

An inverse problem in the economic theory of demand

Problème inverse sur la théorie économique de la demande

Ivar Ekeland^a, Ngalla Djitté^{b,*}

^a *Canada Research Chair in Mathematical Economics, University of British Columbia, Canada*

^b *Laboratoire d'Analyse Numérique et d'Informatique, Université Gaston Berger de Saint-Louis, Sénégal*

Received 10 February 2005; received in revised form 12 October 2005; accepted 17 October 2005

Available online 2 December 2005

Abstract

Given an exchange economy consisting of k consumers, there is an associated collective demand function, which is the sum of the individual demand functions. It maps the price system p to a goods bundle $x(p)$. Conversely, given a map $p \rightarrow x(p)$, it is natural to ask whether it is the collective demand function of a market economy. We answer that question in the case when k is less than the number of goods n . The proof relies on finding convex solutions to a strongly nonlinear system of partial differential equations.

© 2005 Elsevier SAS. All rights reserved.

Résumé

La fonction de demande agrégée d'une société composée de k individus résulte de la sommation de k fonctions de demandes individuelles. Elle fait correspondre à un système de prix p , un vecteur de biens $x(p)$. Inversement, étant donnée une fonction $p \rightarrow x(p)$, est-elle une fonction de demande agrégée ? Nous apporterons une réponse à cette question dans le cas où il y a moins de consommateurs que de biens.

© 2005 Elsevier SAS. All rights reserved.

Keywords: Utility function; Aggregate demand function; Inverse problems; Exterior differential systems; Integral manifolds

Mots-clés : Fonction d'utilité ; Demande agrégée ; Problèmes inverses ; Systèmes différentiels extérieurs ; Éléments intégraux ; Variétés intégrales

1. The disaggregation problem

Consider an exchange economy consisting of k consumers. The space of goods is \mathbb{R}^n , and agent i is characterized by a strictly quasi-concave utility function u_i and a wealth c_i . Given a set of non-negative prices $p \in \mathbb{R}_+^n$, agent i chooses its consumption $x_i(p)$ by solving the optimization problem:

* Corresponding author.

E-mail address: ngalla@ceremade.dauphine.fr (N. Djitté).

$$\max_x u_i(x), \quad (1)$$

$$p'x \leq c_i. \quad (2)$$

Assume that, for every p belonging to a convex subset Ω of \mathbb{R}^n , the solution $x_i(p)$ exists, and satisfies $p'x_i(p) = c_i$. This will be the case, for instance, if the u_i are increasing with respect to every argument, and we can take $\Omega = \mathbb{R}^n$. The individual demands $x_i(p)$ are then well-defined on Ω , and adding them up, we obtain the *market demand*:

$$x(p) := \sum_{i=1}^k x_i(p). \quad (3)$$

It obviously satisfies the Walras law:

$$p'x(p) = \sum_{i=1}^k c_i = c. \quad (4)$$

We are interested in the inverse problem: given a map $p \rightarrow x(p)$ from a subset $\Omega \subset \mathbb{R}^n$ into \mathbb{R}^n , an integer k , and numbers w_i , $1 \leq i \leq k$, can one find strictly concave utility functions $u_i(x)$, $1 \leq i \leq k$, such that the decomposition (3) holds, $x_i(p)$ being the solution of problem (1)? An obvious necessary condition is that $x(p)$ should satisfy the Walras law, but is it sufficient?

This is the famous disaggregation problem, which has a long history. In the case when $k \geq n$ (there are more consumers than goods), it was proved by Chiappori and Ekeland in [4] that the Walras law is sufficient. Much earlier, in a celebrated series of papers, Sonnenschein, Mantel and Debreu had investigated the same question in the case of *excess demand*, that is when the linear constraint is of the form $p'x \leq p'\omega_i$, where $\omega_i \in \mathbb{R}^n$ is the initial endowment of agent i . They proved that if $k \geq n$, the Walras law was sufficient. We refer to [13] for a history of the Sonnenschein–Debreu–Mantel results, and to [5] for a recent proof, in the spirit of the present paper.

Much less is known in the case when $k < n$, except in the case $k = 1$, where the group reduces to one person, so that collective demand coincides with individual demand. It is the aim of the present paper to fill that gap in the case of market demand. We do not treat the case of excess demand, which, to our knowledge, still remains open.

The case $k = 1$ is classical. It was first treated by Antonelli [1], whose results went forgotten and were rediscovered by Slutsky [14]. We summarize these results, and sketch part of the proof; the reader will find the full argument in [12], for instance.

Here and later, we denote differentiation by the symbol D , so that Du denotes the Jacobian matrix of the map u :

$$Du = \left(\frac{\partial u^i}{\partial p_j} \right)_{1 \leq i \leq n, 1 \leq j \leq n}$$

and D^2u denotes the matrix of second derivatives of the function u :

$$D^2u = \left(\frac{\partial^2 u}{\partial p_j \partial p_j} \right)_{1 \leq i \leq n, 1 \leq j \leq n}.$$

Proposition 1. *Let Ω be a convex open subset of \mathbb{R}^n , and $x(p)$ be a C^1 map from Ω into \mathbb{R}^n satisfying the Walras law. A necessary and sufficient condition for $x(p)$ to be the individual demand associated with some quasi-concave utility function u is that the restriction of $Dx(p)$ to $[x(p)]^\perp$ is symmetric and negative definite for every $p \in \Omega$.*

In other words, we can write:

$$Dx = Q + \xi x'$$

where Q is a symmetric, negative definite matrix, and ξ is a vector (so that the last term in the above equation is a matrix of rank 1). This is known in the literature as the *Slutsky condition*. For an analogue of this condition in the case of individual excess demand, we refer to [5].

Let us sketch the proof that the Slutsky condition is necessary. Assume that the utility function u of the sole agent is C^2 , and that the restriction of $D^2u(x(p))$ to $[p]^\perp$ is negative definite. Then the individual demand $x(p)$ is characterized by the first-order optimality condition:

$$\begin{aligned} Du(x(p)) &= \lambda(p)p, \\ p'x(p) &= c \end{aligned}$$

where $\lambda(p) > 0$ is the Lagrange multiplier. Applying the Inverse Function Theorem to this system, one finds that the map $x(p)$ and the function $\lambda(p)$ are well-defined and C^1 on Ω .

Introduce now the indirect demand function:

$$v(p) = \max_x \{u(x) \mid p'x \leq c\}. \quad (5)$$

The function $v(p)$ is quasi-convex and C^2 . More precisely, the restriction of $D^2v(p)$ to $[x(p)]^\perp$ is positive definite. In addition, $u(x)$ can be obtained from $v(p)$ by the formula:

$$u(x) = \min_p \{v(p) \mid p'x \geq c\}. \quad (6)$$

Let us differentiate $v(p)$ by applying the envelope theorem to the formula $v(p) = \max_x \{u(x) - \lambda(p)(c - p'x)\}$. We get:

$$Dv(p) = -\lambda(p)x(p), \quad (7)$$

$$x(p) = -\mu(p)Dv(p) \quad (8)$$

where $\mu(p) = 1/\lambda(p)$. Differentiating, we get:

$$Dx = -\mu D^2v + (D\mu)(Dv)'.$$

The last term on the right-hand side vanish on $[x(p)]^\perp = [Dv(p)]^\perp$, so that the restriction of $Dx(p)$ to $[x(p)]^\perp$ is negative definite, and the Slutsky condition is indeed necessary. The fact that it is also sufficient is proved by using formula (6); we refer to [12] for details.

In the case $k > 1$, there is a natural extension of the Slutsky condition, which was first described by Diewert [6] and by Geanakoplos and Polemarchakis [9] in the context of market demand:

Definition 2. Let Ω be a convex open subset of \mathbb{R}^n , and $x(p)$ be a C^2 map from Ω into \mathbb{R}^n satisfying the Walras law. We shall say that $x(p)$ satisfies the generalized Slutsky condition for k consumers, abbreviated to GS(k), if for every $p \in \Omega$ we have:

$$Dx(p) = Q(p) + \sum_{i=1}^k \xi_i(p) \eta_i(p)' \quad (9)$$

where $Q(p)$ is symmetric and negative definite, and the vectors η_i , $1 \leq i \leq k$, contain $x(p)$ in their linear span.

Proposition 3. Let Ω be a convex open subset of \mathbb{R}^n , and $x(p)$ be a C^2 map from Ω into \mathbb{R}^n satisfying the Walras law. A necessary condition for $x(p)$ to be the market demand for an exchange economy with k consumers is that if $x(p)$ satisfies GS(k) on Ω .

Proof. Rewrite formula (3), using formula (8) for each individual demand $x_i(p)$. We get:

$$x(p) = - \sum_{i=1}^k \mu_i(p) Dv_i(p). \quad (10)$$

Now we differentiate:

$$Dx = - \sum_{i=1}^k \mu_i D^2v_i - \sum_{i=1}^k D\mu_i (Dv_i)'. \quad (11)$$

The first term on the right is a symmetric matrix, and the second has the desired form $\sum_{i=1}^k \xi_i \eta_i'$, with $\eta_i = Dv_i$. Because of relation (10), $x(p)$ belongs to the k -dimensional space $E_k(p) = \text{Span}[(Dv_i(p)) \mid 1 \leq i \leq k]$. Finally,

$E_k(p)^\perp$ is the intersection of all the $[Dv_i(p)]^\perp = [x_i(p)]^\perp$, and the restriction of $D^2v_i(p)$ to $[x_i(p)]^\perp$ is positive definite because of the Slutsky condition. So the restriction of $Dx(p)$ to $E_k(p)^\perp$ must be negative definite. Writing:

$$Dx = \left(- \sum_{i=1}^k \mu_i D^2v_i - \sum_{i=1}^k \alpha_i Dv_i (Dv_i)' \right) - \sum_{i=1}^k (D\mu_i + \alpha_i Dv_i) (Dv_i)'$$

and taking $\alpha_i > 0$ large enough, we find that the first term on the right-hand side is symmetric and negative definite, and this is the GS(k) condition. \square

This paper aims at showing that the GS(k) condition is sufficient. We will prove the following:

Theorem 4. *Let $x(p)$ be an analytic map from some convex neighbourhood Ω_0 of $\bar{p} \in \mathbb{R}^n$ into \mathbb{R}^n satisfying $p'x(p) = w$ for some $w > 0$. Assume that $x(p)$ satisfies the GS(k) condition on Ω_0 for some $k > 1$. Then, given any family $w_i > 0$, $1 \leq i \leq k$, such that $\sum w_i = w$, there exists some convex neighbourhood $\Omega_1 \subset \Omega_0$ of \bar{p} , such that $x(p)$ is the market demand function for an exchange economy with k consumers with wealths w_1, \dots, w_k .*

This theorem will be a consequence of another one, which we state in the next section.

2. A system of nonlinear PDEs

Let us rephrase the disaggregation problem as a system of partial differential equations. We are given a constant c , and a map $x(p)$ satisfying $p'x(p) = c > 0$. We choose positive constants c_1, \dots, c_k which sum up to c , and we seek maps $x_1(p), \dots, x_k(p)$ such that $p'x_i(p) = c_i$ and $x_i(p)$ solves the optimization problem (1), (2) for some strictly quasi-concave function u^i . Substituting the expression (8) for each $x_i(p)$, we get the system of equations:

$$x(p) = - \sum_{i=1}^k \mu_i(p) Dv_i(p), \quad (12)$$

$$p'Dv_i(p) = - \frac{c_i}{\mu_i(p)}, \quad 1 \leq i \leq k. \quad (13)$$

Conversely, if there is a set of functions $v_i(p)$ and $\mu_i(p)$, $1 \leq i \leq k$, satisfying this system, if the $v_i(p)$ are C^2 with D^2v_i positive definite, and if the $\mu_i(p)$ are positive, then the $u_i(x)$ defined by formula (6) will be C^2 and strictly quasi-concave, and $x_i(p)$ will maximize $u_i(x)$ under the constraint $p'x \leq w_i$. So the disaggregation problem is equivalent to finding solutions (v_i, μ_i) , $1 \leq i \leq k$, of the system (12), (13) with the v_i convex and the μ_i positive.

Let us rewrite the system as follows:

$$- \sum_{i=1}^k \mu_i \frac{\partial v_i}{\partial p_j} = x^j(p), \quad 1 \leq j \leq n, \quad (14)$$

$$- \mu_i \sum_{j=1}^n p_j \frac{\partial v_i}{\partial p_j} = c_i, \quad 1 \leq i \leq k. \quad (15)$$

Theorem 5. *Let $x(p)$ be an analytic map from some convex neighbourhood Ω_0 of $\bar{p} \in \mathbb{R}^n$ into \mathbb{R}^n satisfying $p'x(p) = \sum_{i=1}^k c_i$. If $x(p)$ satisfies the GS(k) condition on Ω_0 for some $k > 1$, then there exists some neighbourhood $\Omega_1 \subset \Omega_0$ of \bar{p} , and analytic functions $v_i(p)$ and $\mu_i(p)$, $1 \leq i \leq k$, defined on Ω_1 , with D^2v_i positive definite and $\mu_i > 0$, satisfying equations (14) and (15).*

Of course, we can eliminate the μ_i from the equations. We then get a system of n nonlinear partial differential equations for the k unknown functions v_i :

$$\sum_{i=1}^k \frac{c_i}{\sum_{j=1}^n p_j \frac{\partial v_i}{\partial p_j}} \frac{\partial v_i}{\partial p_j} = x^j(p), \quad 1 \leq j \leq n. \quad (16)$$

This can be written in a more transparent way by introducing the vector field:

$$\pi(p) = \sum_{j=1}^n p_j \frac{\partial}{\partial p_j} \quad (17)$$

and recognizing in the denominator of (16) the derivative of v_i in the direction π , which we denote by $\partial/\partial\pi$:

$$\sum_{i=1}^k c_i \frac{\partial v_i / \partial p_j}{\partial v_i / \partial \pi} = x^j(p), \quad 1 \leq j \leq n. \quad (18)$$

Again, the problem consists in finding convex solutions v_1, \dots, v_k of system (18), and the answer is provided by Theorem 5: it is possible, provided the $x^i(p)$ satisfy the Walras law and the GS(k) condition, and they are analytic.

Before we proceed with the proof, let us comment on related results. If $k < n$, the system is clearly overdetermined (there are more equations than unknown functions), and some compatibility condition on the right-hand sides $x^j(p)$, $1 \leq i \leq n$, is needed in order to find solutions; this is what Theorem 5 provides, with the added twist that we want the solutions to be convex.

When there is a single equation, $k = 1$, the results of the preceding section show that there is a quasi-convex solution $v_1(p) = v(p)$ provided the $x^j(p)$ satisfy the Walras law and the Slutsky condition. This solution is defined globally (there is no need to restrict the initial neighbourhood), and it does not require that the $x^j(p)$ be analytic: C^2 is enough. On the other hand, it does not give a convex solution, only a quasi-convex one, so the conclusion might seem to be weaker. This is not so; indeed, given a quasi-convex function v and a point \bar{p} , we can always find a function $\varphi(t)$ such that $\varphi \circ v$ is convex in some neighbourhood of \bar{p} , and we then take advantage of the following remark.

Lemma 6. *If (v_1, \dots, v_k) is a solution of system (18) near \bar{p} and the $\varphi_i: \mathbb{R} \rightarrow \mathbb{R}$ satisfies $\varphi'(v_i(\bar{p})) \neq 0$, then $(\varphi_1 \circ v_1, \dots, \varphi_k \circ v_k)$ is also a solution of system (18) near \bar{p} .*

Proof. Clear. \square

When $k \geq n$, then the GS(k) condition is vacuous, and all we are left with is the Walras law. Indeed, it has been proved by Chiappori and Ekeland [4] that in that case, the system (14), (15) always has local solutions v_i and μ_i , with the v_i convex and the μ_i positive, provided only that the $x^i(p)$ are analytic and satisfy $\sum p_j x^j(p) = \sum c_i$.

When $k < n$, and we consider only part of the system, that is equations (14) but not the constraints (15), then it has been proved by Ekeland and Nirenberg [8] that there are local solutions v_i and μ_i , with the v_i convex and the μ_i positive, provided the $x^j(p)$ are C^2 (not necessarily analytic) and satisfy the GS(k) condition.

This paper is the first one to consider the full problem with $k < n$. The proof is provided in the next section. It relies, as [4], on the Cartan–Kähler theorem, which is a very sophisticated tool for solving systems of partial differential equations in the analytic framework; see [3] for a detailed statement and proof. Let us conclude by remarking on the requirement that the right-hand sides $x^j(p)$ be analytic. This means that they can be expanded in convergent power series in the neighbourhood of any point p . This is much stronger than simply being indefinitely differentiable: the Taylor series has to converge. The reason this is needed is that the proof of the Cartan–Kähler theorem goes by expanding both sides of the equations in power series and matching coefficients. Whether Theorem 5 still holds true if the right-hand sides $x^j(p)$ are only assumed to be C^∞ is not known; we have investigated the question without success.

3. Solving the system

3.1. Some simplifications

We will give the proof only in the case $k = 2$. This will considerably simplify the notations, and make the argument more transparent by getting rid of half the indices. We will also take $c_1 = c_2 = 1$. Let us restate the problem in that case.

We are given an analytic map $x(p)$ of $\Omega_0 \subset \mathbb{R}^n$ into \mathbb{R}^n satisfying the Walras law:

$$p'x(p) = 2 \quad (19)$$

and the $GS(k)$ condition:

$$Dx(p) = Q(p) + \xi_1(p)\eta_1(p)' + \xi_2(p)\eta_2(p)' \quad (20)$$

where $Q(p)$ is symmetric and negative definite, and $x(p)$ belongs to the 2-dimensional subspace:

$$E_2(p) = \text{Span}[\eta_1(p)\eta_2(p)].$$

We want to find functions $u(p)$, $v(p)$, $\lambda(p)$ and $\mu(p)$ such that u and v are convex, λ and μ are negative, and:

$$\lambda \frac{\partial u}{\partial p_j} + \mu \frac{\partial v}{\partial p_j} = x^j(p), \quad 1 \leq j \leq n, \quad (21)$$

$$\lambda \sum_{j=1}^n p_j \frac{\partial u}{\partial p_j} = 1, \quad (22)$$

$$\mu \sum_{j=1}^n p_j \frac{\partial v}{\partial p_j} = 1. \quad (23)$$

This system naturally splits in two parts: the n Eqs. (21), which express the vector $x(p)$ as a linear combination of the two gradients Du and Dv :

$$\lambda(p)Du + \mu(p)Dv = x(p) \quad (24)$$

and the two Eqs. (22) and (23), which are constraints on the coefficients λ and μ :

$$\lambda(p'Du) = 1, \quad (25)$$

$$\mu(p'Dv) = 1. \quad (26)$$

If we define new variables u^i and v^j by:

$$u^i := \frac{\partial u}{\partial p_i},$$

$$v^i := \frac{\partial v}{\partial p_i}$$

Eqs. (21), (22) and (23) become algebraic relations between the u^i , v^j , λ and μ :

$$\lambda u^i + \mu v^i = x^i(p), \quad 1 \leq i \leq n, \quad (27)$$

$$\lambda \sum_{j=1}^n p_j u^j = 1, \quad (28)$$

$$\mu \sum_{j=1}^n p_j v^j = 1. \quad (29)$$

Define \mathcal{M} to be the set of all $(p_i, u^j, v^k, \lambda, \mu)$, $1 \leq i, j, k \leq n$, satisfying (27), (28) and (29). The problem consists in finding convex functions u and v , negative functions λ and μ such that:

$$(p, Du(p), Dv(p), \lambda(p), \mu(p)) \in \mathcal{M} \quad \forall p. \quad (30)$$

In other words, we are looking for holonomic sections of \mathcal{M} over \mathbb{R}^n .

3.2. Finding general solutions

In order to study the manifold \mathcal{M} and to have a convenient setting for the Cartan–Kähler theorem we will change coordinates in \mathbb{R}^n . We will then apply the Cartan–Kähler theorem to find holonomic sections of \mathcal{M} over \mathbb{R}^n , that is, to find functions u , v , λ and μ which solve the system (21), (22) and (23) but which do not necessarily satisfy the further requirements of convexity and negativity.

Define a differential 1-form ω by:

$$\omega = \sum_{i=1}^n x^i(p) dp_i. \quad (31)$$

Because of the GS(k) condition (20), we have:

$$\omega \wedge d\omega \wedge d\omega = 0. \quad (32)$$

By the Darboux theorem (see [3] or [8]), we can find functions $(q_1(p), q_2(p), q_3(p), q_4(p))$ such that:

$$\omega = q_1 dq_2 + q_3 dq_4. \quad (33)$$

Find other functions $q_i(p)$, $5 \leq i \leq n$, such that the $Dq_i(\bar{p})$, $1 \leq i \leq n$, are linearly independent, and use the q_i as a nonlinear coordinate system near \bar{p} .

Note that, setting $h = q_1/q_3$, we have:

$$\omega \wedge d\omega = (q_3 dq_1 - q_1 dq_3) \wedge dq_2 \wedge dq_4 = q_3^2 dh \wedge dq_2 \wedge dq_4. \quad (34)$$

If there is another splitting of ω , namely $\omega = \lambda du + \mu dv$, then we must also have:

$$\omega \wedge d\omega = (\mu d\lambda - \lambda d\mu) \wedge du \wedge dv = \mu^2 d\left(\frac{\lambda}{\mu}\right) \wedge du \wedge dv. \quad (35)$$

Comparing formulas (34) and (35), we find that (dh, dq_2, dq_4) and $(d(\frac{\lambda}{\mu}), du, dv)$ span the same subspace. It follows that du and dv must belong to the 3-dimensional subspace $E_3(p) = \text{Span}[dq_2, dq_4, dh]$. Writing:

$$du = \sum_{i=1}^n u^i dq_i \quad (36)$$

we find that in fact:

$$du = u^2 dq_2 + u^4 dq_4 + u^h dh = u^2 dq_2 + u^4 dq_4 + \frac{1}{q_3} u^h dq_1 - \frac{q_1}{q_3^2} u^h dq_3. \quad (37)$$

Comparing formulas (36) and (37), we find that:

$$\begin{aligned} q_1 u^1 + q_3 u^3 &= 0, \\ u^i &= 0, \quad 5 \leq i \leq n \end{aligned}$$

and similar relations for the v^j .

In this coordinate system, \mathcal{M} is the set of $(q_i, u^j, v^k, \lambda, \mu)$, $1 \leq i, j, k \leq n$, such that:

$$\lambda u^1 + \mu v^1 = 0, \quad (38)$$

$$\lambda u^2 + \mu v^2 = q_1, \quad (39)$$

$$\lambda u^4 + \mu v^4 = q_3, \quad (40)$$

$$\lambda u^i + \mu v^i = 0, \quad 5 \leq i \leq n, \quad (41)$$

$$u^1 q_1 + u^3 q_3 = 0, \quad (42)$$

$$v^1 q_1 + v^3 q_3 = 0, \quad (43)$$

$$\lambda \sum_{i=1}^4 u^i P_i(q) = 1 \quad (44)$$

where the $P_i(q)$ are the q_i -coordinates of the vector field π , defined by formula (17) in the p_i -coordinates.

All these equations are independent, so that \mathcal{M} is a submanifold of codimension $n + 2$ in \mathbb{R}^{3n+2} , and hence dimension $2n$.

Note that all points in \mathcal{M} satisfy additional relations. From (42) and (43) we deduce that $u^1/v^1 = u^3/v^3$, and equation (38) then implies that:

$$\lambda u^3 + \mu v^3 = 0.$$

Remembering the Walras law, we also find that Eq. (44) implies a similar relation for μ :

$$\mu \sum_{i=1}^4 v^i P_i = \left(\mu \sum_{i=1}^4 v^i P_i + \lambda \sum_{i=1}^4 u^i P_i \right) - \lambda \sum_{i=1}^4 u^i P_i = 1.$$

Consider the exterior differential system on \mathcal{M} :

$$\sum_{i=1}^4 du^i \wedge dq_i = 0, \quad (45)$$

$$\sum_{i=1}^4 dv^i \wedge dq_i = 0, \quad (46)$$

$$dq_1 \wedge \cdots \wedge dq_n \neq 0, \quad (47)$$

where the functions $P_i(q^1, \dots, q^n)$, $1 \leq i \leq n$, are given.

Lemma 7. *Any integral manifold of this system is the graph of a map:*

$$q \rightarrow \left(\frac{\partial u}{\partial q_i}, \frac{\partial v}{\partial q_j}, \lambda, \mu \right)$$

where the functions u , v , λ , μ satisfy Eqs. (21)–(23) in the p_i -coordinates.

Proof. Condition (45) mean that the functions u^i do not depend on the q_j , $5 \leq j$, and that the cross-derivatives $\partial u^i / \partial q_j$ and $\partial u^j / \partial q_i$ are equal, for $1 \leq i, j \leq 4$. By the Poincaré lemma, there is a function $u(q_1, q_2, q_3, q_4)$ such that $u^i = \partial u / \partial q_i$. Similarly, condition (46) means that there is a function $v(q_1, q_2, q_3, q_4)$ such that $v^i = \partial v / \partial q_i$. These partial derivatives lie in \mathcal{M} for all q . Going back to the p_i -coordinates, this means precisely that they satisfy all the equations (21)–(23). \square

We claim that the Cartan–Kähler theorem applies, so that there is an integral manifold going through any given integral element. To prove it, we will have to check successively that the system is closed, that there is an integral element through every point in \mathcal{M} , and that the Cartan test is satisfied.

It is obvious from the form of Eqs. (45) and (46) that the system is closed.

Let us now look for integral elements. Setting:

$$du^i = \sum_{j=1}^n U^{ij} dq_j,$$

$$dv^i = \sum_{j=1}^n V^{ij} dq_j,$$

$$d\lambda = \sum_{j=1}^n L^j dq_j,$$

$$d\mu = \sum_{j=1}^n M^j dx_j$$

and substituting in Eqs. (45) and (46) we get:

$$U^{ij} - U^{ji} = 0, \quad 1 \leq i, j \leq 4, \quad i \neq j, \quad (48)$$

$$V^{ij} - V^{ji} = 0, \quad 1 \leq i, j \leq 4, \quad i \neq j, \quad (49)$$

$$U^{ij} = 0, \quad 1 \leq i \leq 4; \quad 5 \leq j, \quad (50)$$

$$V^{ij} = 0, \quad 1 \leq i \leq 4; \quad 5 \leq j. \quad (51)$$

Note that the U^{ij} , the V^{ij} , the L^j and the M^j also have to satisfy the relations obtained by differentiating Eqs. (38)–(44), which express that we are working in the Grassmannian bundle of n -planes in $T^*\mathcal{M}$. All these equations are linearly independent. Eqs. (48)–(51) number $12 + 4(n-4) + 4(n-4) = 8n - 20$, so that the set of integral elements has codimension $c = 8n - 20$ in the Grassmannian.

Let us now perform Cartan's test. Write:

$$\alpha^i = du^i - \sum_{j=1}^4 U^{ij} dq_j,$$

$$\beta^i = dv^i - \sum_{j=1}^4 V^{ij} dq_j.$$

Note that, because of relations (48) to (51), we have

$$\sum \alpha^i \wedge dq_i = \sum du^i \wedge dq_i = 0,$$

$$\sum \beta^i \wedge dq_i = \sum dv^i \wedge dq_i = 0.$$

We then apply the Cartan procedure, as described in [3]. We have:

$$\begin{aligned} H_0 &= \{0\}, \\ H_1 &= \text{span}[\alpha^1, \beta^1], \\ H_2 &= \text{span}[\alpha^1, \beta^1, \alpha^2, \beta^2], \\ H_3 &= \text{span}[\alpha^1, \beta^1, \alpha^2, \beta^2, \alpha^3, \beta^3], \\ H_i &= \text{span}[\alpha^1, \beta^1, \alpha^2, \beta^2, \alpha^3, \beta^3, \alpha^4, \beta^4] \quad \text{for } 4 \leq i \end{aligned}$$

and hence:

$$c_0 = 0, \quad c_1 = 2, \quad c_2 = 4, \quad c_3 = 6, \quad c_4 = \dots = c_{n-1} = 8,$$

$$C = c_0 + \dots + c_{n-1} = 0 + 2 + 4 + 6 + 8(n-4) = 8n - 20$$

which is exactly the codimension we found earlier: $C = c$. So the exterior differential system passes the Cartan test. By the Cartan–Kähler theorem, for every point $(q_i, u^j, v^k, \lambda, \mu)$, $1 \leq i, j, k \leq n$ in \mathcal{M} , and every integral element $(U^{ij}, V^{k\ell}, L^r, M^s)$ through that point, there is an integral manifold of the exterior differential system (45)–(47).

Of course, the conclusion is coordinate-independent. Let us revert to the original coordinates in \mathbb{R}^n , so that ω no longer has the special form $q_1 dq_2 + q_3 dq_4$. The relations defining an integral element are then obtained directly by writing that they are tangent to holonomic sections of \mathcal{M} . A point in \mathcal{M} is then specified by coordinates $(p_i, u^j, v^k, \lambda, \mu)$, $1 \leq i, j, k \leq n$ satisfying relations (27), (28) and (29). Introduce the notations:

$$\xi = (u^1, \dots, u^n), \quad \eta = (v^1, \dots, v^n).$$

An integral element then is simply a set (U, V, L, M) , where U and V are symmetric $n \times n$ matrices, and L and M are n -vectors satisfying:

$$\Omega = \lambda U + \mu V + L\xi' + M\eta', \tag{52}$$

$$\lambda Up = -(\xi' p)L - \lambda \xi, \tag{53}$$

$$\mu Vp = -(\eta' p)M - \mu \eta. \tag{54}$$

Here Ω is the Jacobian matrix $\partial x^i / \partial p_j$. Note for future use that, by differentiating the relation $\sum x^i(p) p_i = 2$ with respect to p , we get

$$\sum_{i=1}^n \frac{\partial x^i}{\partial p_j} p_i + x^j = 0,$$

so that:

$$\Omega p + \lambda \xi + \mu \eta = 0. \quad (55)$$

The Cartan–Kähler theorem tells us that, given such an integral element at $(p, \xi, \eta, \lambda, \mu)$, there are functions (u, v, λ, μ) such that:

$$D^2 u(\bar{p}) = U, \quad D^2 v(\bar{p}) = V, \quad (56)$$

$$Du(\bar{p}) = \xi, \quad Dv(\bar{p}) = \eta. \quad (57)$$

3.3. Finding convex solutions

Given relations (56) and (57), all we have to prove is that there is an integral element (U, V, L, M) at \bar{p} with U and V positive definite. Then $D^2 u(\bar{p})$ and $D^2 v(\bar{p})$ will be positive definite, and so u and v will be strictly convex in a neighbourhood of \bar{p} .

To do this, we shall use a lemma of Pierre-Louis Lions [11]:

Lemma 8. *Given a symmetric, positive definite matrix Q , a vector p and two vectors q_1 and q_2 satisfying $q_1 + q_2 = Qp$, a necessary and sufficient condition for the existence of two symmetric, positive definite matrices Q_1 and Q_2 such that:*

$$Q = Q_1 + Q_2,$$

$$Q_1 p = q_1,$$

$$Q_2 p = q_2$$

is that:

$$(p, q_1) > 0, \quad (58)$$

$$(p, q_2) > 0, \quad (59)$$

$$(q_1, Q^{-1} q_2) > 0. \quad (60)$$

Proof. Let us set $x = Q^{1/2} p$, $y = Q^{-1/2} q_1$ and $z = Q^{-1/2} q_2$. Set also

$$M = Q^{-1/2} Q_1 Q^{-1/2} \quad \text{and} \quad N = Q^{-1/2} Q_2 Q^{-1/2}.$$

We have:

$$M + N = I, \quad (61)$$

$$Mx = y, \quad (62)$$

$$Nx = z \quad (63)$$

and $y + z = x$. So it is enough to prove the lemma for the particular case where $Q = I$. Let us do it.

Using the three equations, we find easily:

$$(x, y) = (x, Mx) > 0, \quad (64)$$

$$(x, z) = (x, Nx) > 0. \quad (65)$$

Both M and $I - M$ are positive definite. This means that $0 < M < I$ in the sense of symmetric matrices, so that $M^2 < M$. It follows that:

$$(y, z) = (Mx, Nx) = (Mx, (I - M)x) = (x, Mx) - (x, M^2 x) > 0. \quad (66)$$

So the conditions (58), (59), and (60) are necessary.

Let us now prove that they are sufficient. Given three vectors x , y , z satisfying (64), (65), and (66) we seek a symmetric matrix M such that $Mx = y$ and $0 < M < I$. Indeed, we then set $N = I - M$, which is still symmetric and positive definite, and we have $Nx = x - Mx = x - y = z$, as desired.

It is enough to solve the problem in the plane spanned by x and y . Without loss of generality, we assume that $\|x\| = 1$. Take an orthonormal basis (\bar{e}_1, \bar{e}_2) in that plane with $x = (1, 0)$ and $y = (y_1, y_2)$. Eqs. (64), (65) and (66) yield $0 < y_1 < 1$ and $y_1 - \|y\|^2 > 0$.

The matrix we are looking for is:

$$M = \begin{pmatrix} y_1 & y_2 \\ y_2 & c \end{pmatrix}$$

where we will adjust the constant c . Since (x, y) and (x, z) are positive, we have $0 < y_1 < 1$.

Clearly $Mx = y$. We will adjust the constant c to have $0 < M < I$. This requires that the determinant and the trace of M and $(I - M)$ are positive.

Writing that both determinants are positive, we get:

$$\frac{y_2^2}{y_1} < c < 1 - \frac{y_2^2}{1 - y_1}.$$

Since $y_1 > \|y\|^2$, the left-hand side is smaller than the right-hand side, so we can find some c in that interval. Since $0 < y_1 < 1$, we must have $0 < c < 1$.

Writing that both traces are positive, we get:

$$-y_1 < c < 2 - y_1.$$

Since $0 < y_1 < 1$, this inequality follows from $0 < c < 1$. So any c that satisfies the first inequality satisfies the second. \square

We now return to our main argument. By assumption, the $GS(k)$ condition is satisfied, so that there is a symmetric, positive definite $n \times n$ matrix Q and n -vectors L and M such that:

$$\Omega = Q + L\xi' + M\eta'. \quad (67)$$

Lemma 9. Denote by L_0 and M_0 the projections of L and M on $[\xi, \eta]^\perp$ and by Q_0 the restriction of Q to $[\xi, \eta]^\perp$. Without loss of generality, it can be assumed that

$$(\xi'p)(\eta'p)L_0'Q_0^{-1}M_0 > 0.$$

Proof. Take any numbers α , β , γ and δ such that $\beta\gamma - \alpha\delta = 1$. Rewrite Ω as follows:

$$\Omega = Q + \bar{L}\bar{\xi}' + \bar{M}\bar{\eta}'$$

with:

$$\begin{aligned} \bar{L} &= \alpha L + \beta M, & \bar{M} &= \gamma L + \delta M, \\ \bar{\xi} &= -\delta\xi + \gamma\eta, & \bar{\eta} &= \beta\xi - \alpha\eta. \end{aligned}$$

We then have:

$$\begin{aligned} (\bar{\xi}'p)(\bar{\eta}'p)\bar{L}_0'Q_0^{-1}\bar{M}_0 &= [-\delta(\xi'p) + \gamma(\eta'p)][\beta(\xi'p) - \alpha(\eta'p)](\alpha L_0' + \beta M_0')Q_0^{-1}(\gamma L_0 + \delta M_0) \\ &= [-\delta(\xi'p) + \gamma(\eta'p)][\beta(\xi'p) - \alpha(\eta'p)][\alpha\gamma L_0'Q_0^{-1}L_0 + (\alpha\delta + \beta\gamma)L_0'Q_0^{-1}M_0 + \beta\delta M_0'Q_0^{-1}M_0]. \end{aligned}$$

Set the value of δ to $\bar{\delta} = (\eta'p)$. The first term on the right-hand side then changes sign for $\gamma = \bar{\gamma} = (\xi'p)$. We will find values $\bar{\alpha}$ and $\bar{\beta}$, with $\bar{\gamma}\bar{\beta} - \bar{\delta}\bar{\alpha} = 1$ such that the two other terms are non-zero for $(\bar{\alpha}, \bar{\beta}, \bar{\gamma}, \bar{\delta})$. It then follows that the right hand side has two opposite signs for $(\bar{\alpha}, \bar{\beta}, \bar{\gamma} + \varepsilon, \bar{\delta})$ and for $(\bar{\alpha}, \bar{\beta}, \bar{\gamma} - \varepsilon, \bar{\delta})$, when $\varepsilon > 0$ is small enough, and the result obtains. \square

Rewrite Eq. (67) as follows:

$$\Omega = Q + \alpha\xi\xi' + \beta\eta\eta' + (L - \alpha\xi)\xi' + (M - \beta\eta)\eta' \quad (68)$$

with α and $\beta \geq 0$, so that the matrix $Q + \alpha\xi\xi' + \beta\eta\eta'$ is positive definite. Let us apply the lemma of Pierre-Louis Lions to find positive definite matrices Q_1 and Q_2 such that:

$$\begin{aligned} Q_1 + Q_2 &= Q + \alpha\xi\xi' + \beta\eta\eta', \\ Q_1 p &= -(\xi' p)(L - \alpha\xi) - \lambda\xi, \\ Q_2 p &= -(\eta' p)(M - \beta\eta) - \mu\eta. \end{aligned}$$

We first check that $Q_1 p$ and $Q_2 p$ add up to $Qp + \alpha\xi(\xi' p) + \beta\eta(\eta' p)$. Using (68), and (52), we have:

$$\begin{aligned} Q_1 p + Q_2 p &= -(\xi' p)(L - \alpha\xi) - (\eta' p)(M - \beta\eta) - \lambda\xi - \mu\eta, \\ Qp + \alpha\xi(\xi' p) + \beta\eta(\eta' p) &= \Omega p - (L - \alpha\xi)(\xi' p) - (M - \beta\eta)(\eta' p). \end{aligned}$$

Subtracting, we get:

$$Qp - Q_1 p - Q_2 p = \Omega p + \lambda\xi + \mu\eta$$

and the right-hand side vanishes by relation (55).

The first condition holds, and we proceed to the others. Conditions (58) and (59) become:

$$\begin{aligned} -(\xi' p)[(p' L) - \alpha(\xi' p) - \lambda] &> 0, \\ -(\eta' p)[(p' M) - \beta(\eta' p) - \mu] &> 0 \end{aligned}$$

both of which hold true for $\alpha > 0$ and $\beta > 0$ large enough.

When $\alpha, \beta \rightarrow \infty$, we have, for any ξ and η in \mathbb{R}^n :

$$(\xi, [Q + \alpha\xi\xi' + \beta\eta\eta']^{-1}\eta) \rightarrow (\xi_0, Q_0^{-1}\eta_0)$$

where the subscript 0 denotes the projection on $[\xi, \eta]^\perp$. Applying this with $\xi = -(\xi' p)(L - \alpha\xi) - \lambda\xi$ and $\eta = -(\eta' p)(M - \beta\eta) - \mu\eta$, we find:

$$(\xi_0, Q_0^{-1}\eta_0) = (\xi' p)(\eta' p)L'_0 Q_0^{-1} M_0$$

and the right-hand side is positive by the lemma.

The proof is thus happily concluded in the case $k = 2$. In the case $k > 2$, the same argument goes through, except for the last Lemma, which relies on the fact that M and $(I - M)$ commute. One must then use another characterisation, which is due to Inchakov [10]. Without loss of generality, assume $Q = I$, and consider the quadratic form:

$$(Cz, z) = \sum \frac{(x_n, z)^2}{(x_n, y)} - (z, z).$$

Lemma 10. Assume $(x_n, y) \neq 0$ for all n . Then a necessary and sufficient condition for the existence of positive definite matrices M_n such that $M_n y = x_n$ for all n and $\sum M_n = I$ is that $\sum x_n = y$, $(x_n, y) > 0$ for all n , and $(Cz, z) < 0$ for all z not collinear with y .

A proof of Inchakov's lemma is provided in [7].

4. Conclusion

We would like to conclude with a remark of some economic interest. It has been shown by Browning and Chiapori [2] that condition $GS(k)$ is necessary in the case of household demand – that is, when there are public goods in the economy and externalities between consumers. In other words, if $x(p)$ is the demand function of a household with $k < n$ consumers, then it must satisfy $GS(k)$. But it follows from our result that the same $x(p)$ also is the demand function of an exchange economy, where there are only private goods and no externalities. In other words, households and exchange economies cannot be distinguished by looking at the demand functions only.

References

- [1] G. Antonelli, *Sulla teoria matematica della economia politica*, Pisa, 1886; Translated in: J. Chipman, L. Hurwicz, M. Richter, H. Sonnenschein, et al. (Eds.), *Preferences, Utility and Demand*, Harcourt Brace Jovanovich, 1971.
- [2] W. Browning, P.A. Chiappori, Efficient intra-household allocation: a general characterization and empirical tests, *Econometrica* 66 (6) (1998) 1241–1278.
- [3] R.L. Bryant, S.S. Chern, R.B. Gardner, H.L. Goldschmidt, P.A. Griffiths, *Exterior Differential Systems*, Springer-Verlag, New York, 1991.
- [4] P.A. Chiappori, I. Ekeland, Aggregation and market demand: an exterior differential calculus viewpoint, *Econometrica* 67 (1999) 1435–1458.
- [5] P.A. Chiappori, I. Ekeland, Individual excess demands, *J. Math. Economics*, in press.
- [6] W.E. Diewert, Generalized Slutsky conditions for aggregate consumer demand functions, *J. Economic Theory* 15 (1977) 353–362.
- [7] I. Ekeland, Some applications of the Cartan–Kähler theorem to economic theory, in: Erich Kähler, *Mathematische Werke*, de Gruyter, Berlin, 2003, pp. 767–777.
- [8] I. Ekeland, L. Nirenberg, A convex Darboux theorem, *Methods Appl. Math.* 9 (2002) 329–344.
- [9] J. Geanakoplos, H. Polemarchakis, On the disaggregation of excess demand functions, *Econometrica* (1980) 315–333.
- [10] Inchakov, Personal communication, 2000.
- [11] P.L. Lions, Personal communication, Université Paris-Dauphine, 1998.
- [12] A. Mas-Colell, M. Whinston, J. Green, *Microeconomic Theory*, Oxford University Press, 1995.
- [13] W. Shafer, H. Sonnenschein, Market demand and excess demand functions, in: K. Arrow, M. Intriligator (Eds.), *Handbook of Mathematical Economics*, vol. 2, North-Holland, 1982 (Chapter 14).
- [14] E. Slutsky, *Sulla teoria del bilancio del consumatore*, *Giornale degli Economisti e Rivista di Statistica* 3 (15) (1915) 1–26; English translation in: G. Stigler, K. Boulding (Eds.), *Readings in Price Theory*, Richard D. Irwin, 1952.